

Kaushal RK¹, Kumari R² and Kanwar SS¹

¹Department of Biotechnology, ²Department of Chemistry, Himachal Pradesh University, Summer Hill, Shimla-171 005, India
Telfax: +91-177-2832154; kaushalrajeev2@yahoo.com



Lipases (E.C. 3.1.1.3) have been extensively used to achieve various esterification reactions in water restricted or water free media. In the present study a purified thermotolerant alkalophilic extra-cellular lipase of *Bacillus coagulans* MTCC-6375 has been efficiently immobilized onto a synthetic *poly* (MAc-co-DMA-cl-MBAm) hydrogel by surface absorption; and the bound lipase was used to perform short-chain fatty acid ester synthesis in *n*-alkane(s). The hydrogel bound lipase resulted in approximately 64 mM of isoamyl acetate at 55°C in *n*-heptane under shaking in 15 h when vinyl acetate: isoamyl alcohol was used in a ratio of 100 mM: 100 mM. Addition of a molecular sieve (3 Å x 1.5 mm) to the reaction system at a concentration of 25-500 mg per reaction volume had deleterious effect on the conversion of reactants to isoamyl acetate (69.2 mM). During the repetitive esterification under optimal conditions, the hydrogel bound lipase produced 31.3 mM of ester after 4th cycle of reuse. Use of methanol and 2-propanol (instead of isoamyl alcohol) resulted in 78.2 and 64.9 mM of methyl acetate and 2-propyl acetate, respectively under the optimized conditions in *n*-heptane

Key words: *Bacillus coagulans* MTCC-6375; lipase; hydrogel; immobilization.

Introduction

Immobilized enzymes are used in many commercialized products for higher yields. The lipases (E.C. 3.1.1.3) that constitute a most versatile group of enzymes have been used for performing esterification and trans-esterification reactions in organic solvents. In the last few years, there has been an increasing interest in the use of enzymes for the biosynthesis of molecules in organic media [Gargouri *et al.*, 2002, Castillo *et al.*, 2003 and Kanwar *et al.*, 2006]. Esters of short chain fatty acids and alcohols are known as flavor and fragrance compounds that are used in food, beverage, cosmetic and pharmaceutical industries. Isoamyl acetate occurs naturally in the volatile portion of banana fruit and cocoa bean, and is used in perfumes/fragrances, as a flavoring agent in non-alcoholic beverages, ice cream, candy, baked goods, and chewing gum; a solvent for old oil colors, tannins, nitrocellulose, lacquers, paints, formaldehyde, synthetic resins and waxes, celluloid and camphor; for masking unpleasant odors and to perfume shoe polish. Butyl acetate is often employed in combination with *n*-butanol in paints, because *n*-butanol enhances the resistance to blushing and increases the solvency in many cases. In the present study a synthetic *poly*(MAc-co-DMA-cl-DMAm) hydrogel bound lipase of *Bacillus coagulans* MTCC-6375 has been employed in organic media to synthesize isoamyl acetate and a few other short-chain esters by transesterification.

Material and methods

Microorganism :The *Bacillus coagulans* MTCC-6375 isolate was obtained from Department of Biotechnology, Himachal Pradesh University, Summer Hill, Shimla (India).

Synthesis of hydrogel: The hydrogel *poly* (MAc-co-DMA-cl-MBAm) was prepared by co-polymerizing methacrylic acid (23.6 mM) and dodecyl methacrylate (6.85 mM) in acetone (4 ml), in the presence of a cross-linker *N, N*-methylene bisacrylamide (376 mg) and benzoyl peroxide (30 mg) as an initiator.⁵ The reactants were vigorously stirred and transferred inside an airtight glass vial under vacuum. The vial was sealed and heated at 80°C for 30 minutes in a water bath. Insoluble product (polymer) was separated from the reaction mixture by filtration to remove unreacted soluble constituents. Successive washings with water, isoamyl alcohol and

acetone in that order separated the unreacted compounds trapped in the body of the polymer. The polymer was dried completely in an air-oven at 40°C for 24 h to obtain a xerogel.

Immobilization : Purified lipase (8 ml; 35.6 U; 0.52 mg protein/ml) was incubated with 2 g of pre-swollen *poly* (AAc-co-HPMA-cl-MBAm) hydrogel for 18 h at 4°C. The matrix was pre-swollen in Tris buffer (0.05 M, pH 8.5) for 16 h at 8°C. The lipase-bound matrix was filtered through Whatman No.1 filter paper. The bound lipase activity as well as percent protein binding was determined (by estimating unbound protein in the filtrate). Total unbound protein in filtrate was deducted from total protein incubated with the matrix.

Ester synthesis employing immobilized lipase: A reference curve was calibrated between molar concentration (mM) of vinyl acetate and the corresponding area under the peak (retention time 0.48 min). A sample size of 2 μ l was used for GLC analysis. The GLC (Michro-9100, Netel Chromatographs, Thane, India) was programmed for oven temperature 100°C, FID temperature 160°C and injector temperature 160°C. The assay of vinyl acetate was performed on 15% SE30 Chromo WHP column (2 meter X 1.8 inch) using N₂ as a carrier gas (flow rate 30 ml/min).

Determination of amount of isoamyl acetate synthesized: After the completion of esterification at specified time interval, the reaction mixture was withdrawn (2-5 μ l) and subjected to analysis of residual vinyl acetate. The amount of residual vinyl acetate was subtracted from the original amount of vinyl acetate to determine the amount of vinyl acetate consumed. The amount of ester produced in the transesterification reaction was considered to be equivalent to the amount of vinyl acetate consumed in the transesterification reaction.

Results

Kinetics of synthesis of isoamyl acetate: The effect of reaction time on synthesis of isoamyl acetate using immobilized lipase was studied at a temperature of 55°C in *n*-heptane under shaking condition up to 21 h. The synthesis of the ester was time dependent and a maximum amount of isoamyl acetate (64.5 mM) was produced after 15 h of reaction when isoamyl alcohol and vinyl acetate were used at 100 mM each in *n*-heptane (Fig. 1). Thus in subsequent reaction a reaction time of 21 h at 55°C was considered optimum to perform synthesis of isoamyl acetate using hydrogel-bound lipase.

Effect of temperature on synthesis of isoamyl acetate: The effect of change in the reaction temperature on the synthesis of isoamyl acetate by immobilized lipase was also studied. Maximum synthesis (64.6 mM) of isoamyl acetate was obtained at 55°C after 21 h (Fig. 2). At 65°C, there was a marked decrease (55.2 mM) in the ester synthesis, which might be on account of denaturation of the lipase. At 75°C there was a further decline (50.1 mM) in the amount of ester produced.

Effect of solvent (n-alkane) on the ester synthesis: Use of *n*-pentane, *n*-hexane, *n*-hexadecane or *n*-heptadecane reduced the amount of ester formed under similar conditions (Fig. 3). The use of *n*-heptane, *n*-octane and *n*-nonane resulted in 63.7, 63.2 and 66.2 mM of isoamyl acetate. Thus the maximum conversion of reactants into ester was recorded in *n*-nonane (66.2 mM) at 55°C under shaking after 21 h.

Effect of addition of molecular sieve on transesterification: The transesterification reaction resulted in formation of water as a by-product of the reaction, and its removal using a molecular sieve might enhance the synthesis of ester by pushing the reaction equilibrium in the forward direction. However, when the effect of molecular sieve was studied by adding a molecular sieve (25 to 500 mg per reaction volume), a gradual decline (64.0 to 28.4 mM isoamyl acetate) in the amount of ester formed was noticed (Fig. 4). Thus addition of a molecular sieve had a deleterious effect on the transesterification reaction.

Discussion

Lipase is a ubiquitous activity possessed by most of the organisms. Many bacteria, fungi and yeasts produce extra-cellular lipases. Newer diverse microbial sources of lipases have been reported in the recent years. In the present study an extra-cellular alkaline lipase of *Bacillus coagulans* MTCC-6375 was purified and immobilized on to a *poly* (MAc-co-DMA-cl-MBAm) hydrogel. In the esterification reaction, the hydrogel bound lipase of *B. coagulans* optimally produced isoamyl acetate at 55°C in 15 h. However, further increase in the reaction temperature to 65 or 75°C decreased the esterification rate. The alteration in temperature of reaction mixture might interfere with the porosity, hydrophobic character and diffusion of the reactants and/ or products at the catalytic site of enzyme or hydrogel. At temperature more than 55°C, there was no further increase in the amount of ester synthesized, which might be on account of denaturation of the lipase as well as alteration in the native structure of lipase. Recently, the hydrogel-bound lipase of *Bacillus coagulans* MTCC-6375 has been used for optimal synthesis of ethyl laurate at 65°C in 21 h [Kanwar *et. al.*, 2005a]. Also, a synthetic hydrogel immobilized lipase of *P. aeruginosa* BTS-2 was optimally produced ethyl propionate when acid and alcohol was used in an equimolar (100 mM) ratio in the reaction system [Kanwar *et. al.*, 2005b].

The use of alkanes of C-chain length shorter and longer than *n*-heptane decreased the rate of esterification. When *n*-alkane with a shorter C-chain length was used as a solvent, a gradual decrease in rate of isoamyl acetate synthesis was noticed. It was observed that as the log *P* value of an *n*-alkane increased corresponding to increase in the C-chain length of the alkanes (log *P* value for *n*-pentane, *n*-hexane, *n*-heptane, and *n*-nonane is 3, 3.5, 4 and 5, respectively), the hydrophobicity of the alkane *i.e.* solvent also increased in that order, and that appeared to be very important for modulation of catalytic activity of hydrogel-bound *B. coagulans* lipase. Thus it appeared that choice of an appropriate solvent system was critical for the synthesis of ester using hydrogel-immobilized lipase of *Bacillus coagulans*. A relatively higher hydrophobicity appeared to be an important property to be built into a synthetic hydrogel for increasing the catalytic activity of bound lipase of *B. coagulans*. Higher the log *P*, the more hydrophobic is the solvent. In a recent study using hydrogel-immobilized lipase of *P. aeruginosa* BTS-2, we have shown a similar effect that resulted in an enhanced synthesis of ethyl propionate in *n*-nonane.³⁹ However, previously reported lipase of *P. aeruginosa* was more efficient in synthesis of ester as indicated by approximately 94% conversion of reactants into ester. In another study, hydrogel-bound *B. coagulans* MTCC-6375 lipase could achieve 61% conversion of reactants into ethyl laurate.³⁶ This lower conversion into ester might be due to use of a higher carbon chain length fatty acid (lauric acid, C 12) than the one used previously (propionic acid, C 3). It appeared that *B. coagulans* MTCC-6375 lipase is inherently less efficient in performing esterification than the *Pseudomonas* lipase as in the³ present study approximately 64-67 mM of isoamyl acetate could be produced by manipulating various physical and kinetic parameters.

Conclusion

The present study showed that *poly* (MAc-co-DMA-cl-MBAm) hydrogel-immobilized-lipase of *Bacillus coagulans* efficiently performed esterification of isoamyl alcohol and vinyl acetate into isoamyl acetate at a temperature of 55°C, addition of molecular sieve in conjunction with a hydrophobic matrix caused a decline in the rate of esterification, and an equi-molar ratio of isoamyl alcohol as well as vinyl acetate was necessary for achieving an optimal rate of trans-esterification. Foregoing discussion indicated that a systematic approach could be devised on the basis of solvent engineering to favor optimal synthesis of an ester of interest. Many studies have shown that smaller fatty acids are involved in inhibition of enzyme at higher concentration. Generally, commercial enzymes must be very rugged to remain active for prolonged duration when exposed to an adverse environment. This study is among very few reports on the use of a synthetic hydrogel bound extra-cellular alkaline lipase of a moderately thermotolerant *B. coagulans* for the synthesis of isoamyl acetate and a few-other alcohol-acetate esters, of short chain lengths in a water free organic solvent system. The efficacy of this lipase to catalyze inter-esterification and transesterification

reactions in organic solvents/ water restricted/ controlled media are still required to be explored in future studies.




Fig. 1: Effect of reaction time on synthesis of isoamyl acetate.




Fig. 2: Effect of reaction temperature on synthesis of isoamyl acetate.




Fig.3: Effect of solvent (n-alkane) on synthesis of isoamyl acetate.




Fig. 4: Effect of molecular sieve on synthesis of isoamyl acetate.

References

4

- Gargouri M. et al.(2002), *Synthesis of a novel macrolactone by lipase-catalyzed intra-esterification of hydroxy-fatty acid in organic media.*, J Biotechnol, 92, 259.
- Castillo E et. al.(2003), *Lipase catalyzed synthesis of xylitol monoesters: solvent engineering approach*, J Biotechnol, 102, 251.
- Kanwar S. S. et. al.(2005a), Effect of solvents and kinetic parameters on synthesis of ethyl propionate catalyzed by poly (AAc-co-HPMA-cl-MBAm)-matrix-immobilized lipase of *Pseudomonas aeruginosa* BTS-2. World J Microbiol Biotechnol , 21, 1037.
- Kanwar S. S et al.(2005b), Methods for inhibition of residual lipase activity in colorimetric assay: A comparative study. Indian J Biochem and Biophy, 42, 233.
- Kanwar SS et al.(2006), *Catalytic potential of a poly (AAc-co-HPMA-cl-MBAm)-matrix-immobilized lipase from a thermotolerant Pseudomonas aeruginosa MTCC 4713*, J Appl Polym Sci, 100,4252..